



Nuclear interactions of low-energy cosmic rays with the interstellar medium

V. Tatischeff and J. Kiener

Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse, IN2P3/CNRS and Univ Paris-Sud, 91405 Orsay, France, e-mail: Vincent.Tatischeff@csnsm.in2p3.fr

Abstract. Cosmic rays of kinetic energies below ~ 1 GeV nucleon $^{-1}$ are thought to play a key role in the chemistry and dynamics of the interstellar medium. They are also thought to be responsible for nucleosynthesis of the light elements Li, Be, and B. However, very little is known about the flux and composition of low-energy cosmic rays since the solar modulation effect makes impossible a direct detection of these particles near Earth. We first discuss the information that the light elements have brought to cosmic-ray studies. We then discuss the prospects for detection of nuclear gamma-ray line emission produced by interaction of low-energy cosmic rays with interstellar nuclei.

Key words. Cosmic rays – Nuclear reactions, nucleosynthesis, abundances – Gamma rays: ISM

1. Introduction

Low-energy cosmic rays (LECRs) of kinetic energies $\lesssim 1$ GeV nucleon $^{-1}$ are thought to be a major player in the process of star formation. They are a primary source of ionization of heavily shielded, dense molecular clouds and the resulting ionization fraction conditions both the chemistry in these regions and the coupling of the gas with the ambient magnetic field. LECRs also represent an important source of heating that contribute to hold molecular cores in equilibrium against gravitational forces. In addition, LECRs are thought to drive large-scale magnetohydrodynamic turbulence and cause amplification of magnetic field in the interstellar medium (ISM; see, e.g., Hanasz et al. in these proceedings).

Despite LECRs being a fundamental component of the Galactic ecosystem, their compo-

sition and flux are very uncertain. This is partly because these low-energy particles cannot be detected near Earth due to the solar modulation effect. Although the Voyager 1 and 2 spacecrafts are approaching the heliopause (i.e. the contact discontinuity between the solar wind and the ISM) the twin probes will probably not be able to observe the local interstellar spectrum of CRs below 1 GeV nucleon $^{-1}$, because of the additional modulation likely occurring in the outer heliosheath (Scherer et al. 2011, and Fichtner in these proceedings).

Thus, for still a long time our knowledge of the properties of LECRs will be based mainly on analysis of radiation produced by interaction of these particles with the ISM. In particular, a very promising way to study hadronic CRs (by far the most abundant in the Galaxy) below the kinetic energy threshold for production of neutral pions (280 MeV for $p + p$ collisions) would be to detect characteristic γ -ray




Fig. 1. Be abundances vs. $[O/H]$ ($[O/H] = \log[(O/H)/(O/H)_\odot]$, where $(O/H)/(O/H)_\odot$ is the O abundance relative to its solar value). Data: Rich & Boesgaard (2009); Tan et al. (2009). The solid line shows a fit to the data with a secondary Be production component plus a primary component of constant $Be/O = 1.3 \times 10^{-8}$.

lines produced by nuclear collisions of CRs with interstellar matter (Sect. 3). However, before such a detection will be made possible by a significant improvement of the sensitivity of the γ -ray space instruments, useful information on LECRs can already be obtained from measurements of cosmic abundances of the light elements Li, Be, and B (Sect. 2).

2. Nucleosynthesis of the light elements and origin of cosmic rays

There are two basic reasons to assume that Li, Be, and B are significantly produced by nuclear interaction of galactic cosmic rays (GCRs) with the ISM. First, the abundances of these species relative to, e.g., Si, are much higher in the GCRs than in the solar system, by factors ranging from 10^4 to 10^6 . Second, as first shown by Reeves et al. (1970), the pre-solar abundances of these elements as measured in meteorites can be approximately reproduced by assuming that they are synthesized by fragmentation of interstellar (resp. CR) C, N and O by CR (resp. interstellar) protons and α -particles, plus a contribution from $\alpha + \alpha$ reactions for Li production, and further assuming that the GCR flux and CNO abundances have not changed throughout the history of the

Galaxy¹. Quantitatively, Reeves et al. (1970) found that a flux of CR protons averaged over ~ 10 Gyr $F_p(E > 30 \text{ MeV}) \sim 15 \text{ cm}^{-2} \text{ s}^{-1}$ would be needed to reproduce the pre-solar abundance of Li (sic). In comparison, the various estimates of the local interstellar spectrum (LIS) proposed in the literature over the past few decades give $F_p(E > 30 \text{ MeV})$ in the range $\sim 5 - 40 \text{ cm}^{-2} \text{ s}^{-1}$ (see Indriolo et al. 2009, and references therein).

However, a full calculation of the abundances of the light elements must take into account the CR history and chemical evolution of the Galaxy. In such a framework, it was long expected that Be and B should be produced as secondary species by fragmentation of the increasingly abundant CNO nuclei in both the GCRs and the ISM, that is that Be and B abundances measured in stellar atmospheres should vary *quadratically* with the star metallicity. But observations made from the 1990's reveal that Be and B abundances increase *linearly* with metallicity (or with $[O/H]$, see Fig. 1), at least in old halo stars of low metallicity. The observed B evolution could be accounted for by a predominant production of ^{11}B by ν spallation of ^{12}C in core-collapse supernovae (Woosley et al. 1990), which is a primary production process. But the observed evolution of Be is still not well understood.

2.1. Nucleosynthesis of primary Be in supernova remnants

Synthesis of *primary* Be is expected to occur in every supernova remnant (SNR) producing CRs, because before being released in the ISM at the end of the Sedov-Taylor phase, the particles accelerated at the blast wave can interact within the remnant with SN ejecta enriched in freshly synthesized C and O nuclei (Parizot & Drury 1999a,b). A simple estimate of this Be production can be obtained by assuming that a constant fraction θ_{CR} of the avail-

¹ It is now admitted, however, that only ^6Li , ^9Be , and ^{10}B are pure products of GCR nucleosynthesis, ^7Li being also significantly produced in the Big Bang and in stars and ^{11}B by ν -induced spallation in core-collapse supernovae (see Prantzos 2010).




Fig. 2. Evolution as a function of age of the Be production rate per synthesized O atom (*black curves, left axis*) and the total luminosity in γ -rays > 100 MeV (*red curves, right axis*), for a SNR expanding into a medium of H density n_{ISM} (lower curves: $n_{\text{ISM}}=1 \text{ cm}^{-3}$; upper curves: $n_{\text{ISM}}=100 \text{ cm}^{-3}$).

able mechanical power processed by the blast wave is continuously transformed into kinetic energy of CRs, that can produce Be while being advected with the plasma downstream the shock front. The medium inside the SNR is taken to be well-mixed and thus homogeneous. Neglecting in first approximation the adiabatic and Coulomb energy losses of the fast particles, the Be production rate per O-atom in the downstream plasma can then be estimated to be $(d\text{Be}/dt)/\text{O} = q_{\text{Be}}\epsilon_{\text{CR}}$, where $q_{\text{Be}} \approx 10^{-13} \text{ s}^{-1} (\text{erg cm}^{-3})^{-1} (\text{O-atom})^{-1}$ is the Be production rate normalized to the CR energy density, which post-shock value is $\epsilon_{\text{CR}} = k\theta_{\text{CR}}\rho_{\text{ISM}}V_s^2$. Here, V_s is the forward shock velocity, ρ_{ISM} the mass density of the ISM surrounding the SNR and k a factor of order unity that depends on the equation of state of the shocked gas. We calculated q_{Be} assuming for the CR momentum spectrum a phase-space distribution resulting from diffusive shock acceleration, $f_{\text{CR}}(p) \propto p^{-s}$ with $4 < s < 4.5$, and for the relative abundances of the CNO elements in the SN ejecta C:N:O=0.2:0.05:1 (Woosley & Weaver 1995).

Calculated Be production rates are shown in Fig. 2 as a function of time after the SN explosion. The evolution of V_s is taken from

the blast wave solution of Truelove & McKee (1999), assuming for the supernova energy, the mass of the ejecta, and the power-law index of the outer ejecta density profile: $E_{\text{SN}} = 1.5 \times 10^{51} \text{ erg}$, $M_{\text{ej}} = 10 M_{\odot}$ and $n = 10$, respectively. Also shown in Fig. 2 is the total luminosity of the SNR in γ -rays > 100 MeV, which we calculated from the same simple formalism but using the γ -ray emissivity $q_{\gamma} = 5 \times 10^{-14} \text{ s}^{-1} (\text{erg cm}^{-3})^{-1} (\text{H-atom})^{-1}$ (Drury et al. 1994). We see that both the instantaneous Be production rate and the γ -ray luminosity increase with the density of the ambient ISM, but also that the blast wave evolution is more rapid for larger values of n_{ISM} (e.g. the end of the Sedov phase occurs at $t_{\text{rad}} = 31900$ and 2790 yrs for $n_{\text{ISM}} = 1$ and 100 cm^{-3} , respectively). As a result, the final Be/O abundance ratio obtained by integration of the Be production rate over the SNR lifetime obeys the following scaling relation:

$$\frac{\text{Be}}{\text{O}} \approx 7.1 \times 10^{-10} \left(\frac{\theta_{\text{CR}}}{0.3} \right) \left(\frac{n_{\text{ISM}}}{1 \text{ cm}^{-3}} \right)^{0.64}. \quad (1)$$

Thus, a large ambient medium density $n_{\text{ISM}} \approx 100 \text{ cm}^{-3}$ would be needed to account for the observed abundance ratio in low-metallicity stars, $(\text{Be}/\text{O})_{\text{obs}} \approx 1.3 \times 10^{-8}$ (see Fig. 1), which is not realistic. This robust conclusion is in good agreement with the results of the more detailed calculations of Parizot & Drury (1999a,b). Moreover, as already pointed out by Feltzing & Gustafsson (1994), a present-day production of Be in SNRs at the level of $(\text{Be}/\text{O})_{\text{obs}}$ would be accompanied by fluxes of γ -rays > 100 MeV above those observed from several SNRs and the Galaxy.

2.2. On the origin of cosmic rays

Thus, the observed quasi-linear increase of Be abundance with metallicity implies that a significant amount of CNO nuclei in the GCRs were – and possibly still are – accelerated out of freshly synthesized matter. A first model discussed in the literature is that the GCRs are accelerated mostly out of SN ejecta rather than the average ISM. Thus, Lingenfelter et al. (1998) suggested that high-velocity grains

formed in SN ejecta could be the source of CR metals, which may be accelerated either in the forward or reverse shocks of SNe. But the absence of radioactive ^{59}Ni in the GCRs makes this scenario unlikely, because this observation suggests a delay $\gtrsim 10^5$ years between nucleosynthesis of iron-peak nuclei and their acceleration to CR energies (Wiedenbeck et al. 1999). Indeed ^{59}Ni decays by electron capture with a half life of 7.6×10^4 years for the neutral atoms, but this decay is suppressed once the ^{59}Ni ions are accelerated and fully stripped.

Another scenario is that the CNO isotopes of the GCRs are mainly accelerated from the fresh wind material of massive stars, especially Wolf-Rayet stars, as it is hit by the blast wave of the subsequent SN explosions. Prantzos (2010) pointed out that models of fast rotating massive stars predict an ejection of winds enriched in CNO elements even at low metallicity and suggested that these wind sources could produce essentially constant CNO abundances in the GCRs at all times. According to Ramaty et al. (1997), the measured Be abundances can be explained if each Galactic SN supplies on average a total energy of $\sim 10^{50}$ erg to GCRs of current epoch composition. Assuming a typical CR acceleration efficiency of $\sim 10\%$, it means that the SN blast wave has to process a large fraction of the total explosion energy ($\sim 10^{51}$ erg) while interacting with a medium enriched in freshly synthesized CNO nuclei. In other words, the energetics of Be production implies that, on average per Galactic SN, the mass of the CNO-rich material ahead of the forward shock cannot be much lower than that of the SN ejecta. Models for non-rotating massive stars do not predict a significant ejection of CNO-rich winds at low metallicity. Those for fast-rotating, metal-poor, massive stars do display a substantial wind enriched in He-burning products after the main sequence, but only for stars of initial mass $\geq 60 M_{\odot}$ (Decressin et al. 2007). For the Salpeter's initial mass function, such very massive stars represent no more than $\sim 5\%$ of all stars giving rise to SNe. We also note that the SN explosion of a fast spinning star could favor matter ejection through jets aligned along the rotation axis (Decressin et al. 2007), such that the resulting blast wave

may not interact with all the wind material. For these reasons, it is not clear to us that the scenario recently put forward by Prantzos (2010) can account for enough Be production in the early Galaxy.

A third possibility is that the GCRs are accelerated primarily out of material inside superbubbles of hot gas generated by the winds and SN explosions of massive stars formed in OB associations (Higdon et al. 1998; Parizot & Drury 1999c). The majority of core-collapse SNe likely occur in these cavities. Shock waves propagating in superbubbles can accelerate matter enriched in winds and ejecta of SN explosions of massive stars born in the same OB association, thus producing GCRs of essentially constant metallicity throughout the age of the Galaxy. This model was shown to be compatible with the absence of ^{59}Ni in the GCRs (Binns et al. 2008). However, the superbubble scenario for the origin of the GCRs may suffer from two problems. First, contrary to previous results (e.g., Higdon & Lingenfelter 2003), Prantzos shows in these proceedings that the composition of superbubble gas can probably not account for the well-known anomalous $^{22}\text{Ne}/^{20}\text{Ne}$ abundance ratio in the CR source, which is 5.3 times higher than in the solar system. Moreover, specific CR acceleration effects are expected in superbubbles due to the presence of multiple shocks and plasma turbulence between the shocks (Bykov & Fleishman 1992; Parizot et al. 2004; Ferrand & Marcowith 2010). As a result, the phase-space spectrum of CRs produced in superbubbles is predicted to be harder (resp. softer) than p^{-4} (the "universal" distribution produced by diffuse strong shock acceleration in test-particle regime) below (resp. above) a critical momentum that depends on the superbubble parameters (Ferrand & Marcowith 2010). This prediction is not supported by recent *Fermi* LAT γ -ray observations (see Mizuno et al. 2011) that show the spectrum of the γ -ray-producing CRs to be spatially uniform in the outer Galaxy and consistent with the LIS (see also Tibaldo et al. in these proceedings).

It is possible that the bulk of primary Be is in fact not synthesized by the GCRs themselves, but by a separate LECR component

(Cassé et al. 1995) that could be produced in superbubbles (Vangioni-Flam et al. 1998). These LECRs might be accelerated at early times during the superbubble lifetime, when the dynamics of the system is powered by the strong winds of massive stars. The standard GCRs would then be produced later on, by regular Fermi (i.e. diffusive shock) acceleration in SN blast waves first propagating in the winds of the massive progenitor stars and then in the local ISM (in order to explain the $^{22}\text{Ne}/^{20}\text{Ne}$ abundance ratio in the GCRs, see Prantzos in the proceedings). The postulated LECR component could account for the large ionization rate deduced from H_3^+ abundances in diffuse interstellar clouds (Indriolo et al. 2009), although the extra ionization could also be due to LECR electrons (Padovani et al. 2009).

3. Nuclear gamma-ray line emission

The most compelling evidence for a significant hadronic LECR component would come from the detection of nuclear γ -ray lines produced by interaction of these fast particles with interstellar matter. The strongest lines from the most abundant nuclei are preferentially produced in the CR energy range below a few hundred MeV per nucleon. We estimate here the diffuse Galactic emission in these lines; potential γ -ray line sources such as SNRs will be discussed elsewhere.

Indriolo et al. (2009) proposed several LECR spectra which, added to the standard CR spectrum, would produce the observed mean ionization rate of diffuse molecular clouds $\zeta_{\text{CR}} = 4 \times 10^{-16} \text{ s}^{-1}$. We used a slightly modified version of the broken power-law spectrum of these authors, with the CR composition measured at $\sim 1 \text{ GeV nucleon}^{-1}$ kept constant down to $2 \text{ MeV nucleon}^{-1}$, contrary to Indriolo et al. (2009) who used solar abundances below $200 \text{ MeV nucleon}^{-1}$. This required a slightly changed power-law index of 1.65 for the LECR component to get the same ζ_{CR} . It was verified that the γ -ray emission from π^0 decay for such an interstellar CR spectrum reproduces the local γ -ray emissivity observed by *Fermi* LAT in the range $0.1 - 10 \text{ GeV}$ (Abdo et al. 2009).




Fig. 3. Calculated inner Galaxy nuclear γ -ray line spectrum, with indications of the emitting nuclei for some prominent lines.

For the calculation of the nuclear γ -ray line emission, cross sections for strong lines were taken mainly from Murphy et al. (2009) with some modifications taking into account new nuclear data of Benhabiles et al. (2011). The production of weaker lines and γ -ray emitting radioactive nuclei was calculated with the nuclear reaction code TALYS (Koning et al. 2008). For the interstellar matter, we used solar abundances with 2 times solar metallicity to take into account the metal enrichment of the inner Galaxy. Parallely to the nuclear lines, the γ -ray emission following π^0 production and decay was calculated with the model of Kamae et al. (2006). Adjusting the latter to the pion-decay spectrum of the inner Galaxy at $0.1 - 100 \text{ GeV}$ as deduced from *Fermi* LAT observations (Strong et al. 2011) provided the absolute normalization. The calculated γ -ray line spectrum is shown in Fig. 3. Besides strong narrow lines, the nuclear line emission is composed of broad lines produced by the CR heavy-ion component and of thousands of weaker lines. Adding all up results in a complex and extremely rich emission spectrum.

A very promising feature for an observation with future space-borne γ -ray telescopes is the characteristic bump above $E_\gamma \sim 3 \text{ MeV}$ with several strong lines of ^{12}C and ^{16}O . The calculated flux in the $3 - 10 \text{ MeV}$ band inte-

grated over the inner Galaxy ($300^\circ \leq l \leq 60^\circ$, $|b| \leq 10^\circ$) amounts to $7 \times 10^{-5} \text{ cm}^{-2} \text{ s}^{-1}$. This can be compared with the estimated continuum sensitivities of some recent γ -ray telescope projects, like, e.g., the GRIPS, DUAL and CAPSiTT proposals for the latest (2010) ESA Cosmic Vision 2015–2025 call for a Medium-size space mission. The proposed instruments feature a factor of typically 10 – 100 sensitivity improvement with respect to *CGRO/COMPTEL* and *INTEGRAL/SPI*. Thus, the CAPSiTT 3σ continuum sensitivity at 5 MeV in a 5-year all-sky survey for such a spatially extended emission is estimated to be $\sim 10^{-5} \text{ cm}^{-2} \text{ s}^{-1}$, which would assure a comfortable detection of the predicted nuclear line emission.

4. Conclusions

The observed primary evolution of Be versus [Fe/H] (or [O/H]) shows that some CRs are accelerated out of freshly synthesized matter. The most promising sites of acceleration of these CRs appears to be superbubbles, which, however, are probably not the source of the standard GCRs detected near Earth. But superbubbles may produce a distinct component of LECRs during their early, stellar-wind-dominated phase. This supplementary CR component could account for the large ionization rate of diffuse interstellar clouds.

Nuclear interaction γ -ray lines could provide one of the best ways of studying the origin of LECRs and their role in the Galactic ecosystem. The detection of these lines should be a major objective for the next-generation of MeV γ -ray telescopes.

References

- Abdo, A. A. et al. 2009, *ApJ*, 703, 1249
 Benhabiles-Mezhoud, H. et al. 2011, *Phys. Rev. C*, C 83, 024603
 Binns, W. R., et al. 2008, *New Astron. Rev.*, 52, 427
 Bykov, A. M., & Fleishman, G. D. 1992, *MNRAS*, 255, 269
 Cassé, M., Lehoucq, R., & Vangioni-Flam, E. 1995, *Nature*, 373, 318
 Decressin, T., et al. 2007, *A&A*, 464, 1029
 Drury, L. O., Aharonian, F. A., & Voelk, H. J. 1994, *A&A*, 287, 959
 Feltzing, S., & Gustafsson, B. 1994, *ApJ*, 423, 68
 Ferrand, G., & Marcowith, A. 2010, *A&A*, 510, A101
 Higdon, J. C., Lingenfelter, R. E., & Ramaty, R. 1998, *ApJ*, 509, L33
 Higdon, J. C., & Lingenfelter, R. E. 2003, *ApJ*, 590, 822
 Indriolo, N., Fields, B. D., & McCall, B. J. 2009, *ApJ*, 694, 257
 Kamae, T., et al. 2006, *ApJ*, 647, 692
 Koning, A. J., Hilaire, S., & Duijvestijn, M. C. 2008, *Int. Conf. on Nucl. Data for Science and Tech.*, April 22-27, 2007, Nice, France, eds O. Bersillon et al., EDP Sciences, p. 211
 Lingenfelter, R. E., Ramaty, R., & Kozlovsky, B. 1998, *ApJ*, 500, L153
 Mizuno, T. et al. 2011, *Proc. of the 32nd Int. Cosmic Ray Conf.*, Beijing 2011, in press
 Murphy, R. J., et al. 2009, *ApJS*, 183, 142
 Padovani, M., Galli, D., & Glassgold, A. E. 2009, *A&A*, 501, 619
 Parizot, E., & Drury, L. 1999a, *A&A*, 346, 329
 Parizot, E., & Drury, L. 1999b, *A&A*, 346, 686
 Parizot, E., & Drury, L. 1999c, *A&A*, 349, 673
 Parizot, E., et al. 2004, *A&A*, 424, 747
 Prantzos, N. 2010, *IAU Symposium*, 268, 473
 Ramaty, R., et al. 1997, *ApJ*, 488, 730
 Reeves, H., Fowler, W. A., & Hoyle, F. 1970, *Nature*, 226, 727
 Rich, J. A., & Boesgaard, A. M. 2009, *ApJ*, 701, 1519
 Scherer, K., et al. 2011, *ApJ*, 735, 128
 Strong, A. W., et al. 2011, *ICATPP Conf. on Cosmic Rays for Particle and Astroparticle Physics*, Villa Olmo (Como, Italy), 7-8 October, 2010, in press; *arXiv* 1011.1381
 Tan, K. F., Shi, J. R., & Zhao, G. 2009, *MNRAS*, 392, 205
 Truelove, J. K., & McKee, C. F. 1999, *ApJS*, 120, 299
 Vangioni-Flam, E., et al. 1998, *A&A*, 337, 714
 Wiedenbeck, M. E., et al. 1999, *ApJ*, 523, L61
 Woolsley, S. E., et al. 1990, *ApJ*, 356, 272
 Woolsley, S. E., & Weaver, T. A. 1995, *ApJS*, 101, 181